

## 2018 Continues Record Global Ocean Warming

Lijing CHENG<sup>\*1</sup>, Jiang ZHU<sup>1</sup>, John ABRAHAM<sup>2</sup>, Kevin E. TRENBERTH<sup>3</sup>, John T. FASULLO<sup>3</sup>,  
Bin ZHANG<sup>4,5</sup>, Fujiang YU<sup>6</sup>, Liying WAN<sup>6</sup>, Xingrong CHEN<sup>6</sup>, and Xiangzhou SONG<sup>7,8</sup>

<sup>1</sup>*International Center for Climate and Environment Sciences, Institute of Atmospheric Physics,  
Chinese Academy of Sciences, Beijing 100029, China*

<sup>2</sup>*University of St. Thomas, School of Engineering, Minnesota 55105, USA*

<sup>3</sup>*National Center for Atmospheric Research, Boulder, Colorado 80307, USA*

<sup>4</sup>*Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China*

<sup>5</sup>*Center for Ocean Mega-Science, Chinese Academy of Sciences, Qingdao 266071, China*

<sup>6</sup>*National Marine Environmental Forecasting Center, Ministry of Natural Resources of China, Beijing 100081, China*

<sup>7</sup>*College of Oceanography, Hohai University, Nanjing 210098, China*

<sup>8</sup>*Bureau of Territorial and Spatial Planning, Ministry of Natural Resources, Beijing 100812, China*

(Received 24 December 2018; revised 7 January 2019; accepted 8 January 2019)

**Citation:** Cheng, L. J., and Coauthors, 2019: 2018 continues record global ocean warming. *Adv. Atmos. Sci.*, **36**(3), 249–252, <https://doi.org/10.1007/s00376-019-8276-x>.

The increasing heat-trapping gases emitted by human activities into the atmosphere produce an energy imbalance between incoming solar radiation and outgoing longwave radiation that leads to global heating (Rhein et al., 2013; Trenberth et al., 2014; von Schuckmann et al., 2016). The vast majority of global warming heat ends up deposited in the world's oceans, and ocean heat content (OHC) change is one of the best—if not the best—metric for climate change (Cheng et al., 2019). In 2018, continued record heat was measured in the Earth's climate system. In fact, 2018 has set a new record of ocean heating, surpassing 2017, which was the previous warmest year ever recorded (Cheng et al., 2018) (Fig. 1).


Based on the new update of the Institute of Atmospheric Physics (IAP) ocean analysis (see “Data and methods” section), the total ocean heat anomaly in 2018, relative to a 1981–2010 baseline, and for the upper 2000 m of the world's oceans, is  $(19.67 \pm 0.83) \times 10^{22}$  J. This level of thermal energy places 2018 as the hottest year ever recorded. Figure 1 shows the progression of upper 2000 m OHC since the late 1950s. The ranking of the five warmest years (Table 1) makes the past five years the warmest years on record. This supports the provisional announcement by the World Meteorological Organization in November 2018 that “the ocean heat content was the highest or 2nd highest on record”. And the new record in 2018 confirms the perspective (Cheng et al., 2019) that ocean warming continues and has been accelerating since the 1990s (compared with 1960 to the 1980s).

The heating was distributed throughout the world's oceans, with the vast majority of regions showing an increase in thermal energy (Fig. 1). The Southern Ocean (south of 30°S) and Pacific Ocean showed more warming than the Atlantic Ocean and Indian Ocean (Shi et al., 2018; Swart et al., 2018). The Southern Ocean and Pacific Ocean (north of 30°S) were  $(6.91 \pm 1.70) \times 10^{22}$  and  $(5.97 \pm 1.07) \times 10^{22}$  J above the 1981–2010 period, respectively. In comparison, the values for the Atlantic Ocean (north of 30°S) and Indian Ocean (north of 30°S) were  $(4.95 \pm 1.97) \times 10^{22}$  and  $(1.84 \pm 1.97) \times 10^{22}$  J. The spatial pattern also indicates some imprints of internal variability in the climate system (Cheng et al., 2018). For example, in 2018, the tropical Pacific transformed from a La Niña state in the early part of the year into a weak El Niño condition later (see “Data and methods” section). The heat was recharged in the equatorial regions within 10°S–5°N (Fig. 1), leaving negative OHC anomalies in the north subtropical Pacific Ocean (5°–15°N). The Indian Ocean continued its rapid warming trend since the 1990s, partly linked to both anthropogenic forcing and the Pacific Decadal Oscillation (Li et al., 2018).

Increases in ocean heat are incontrovertible proof that the Earth is warming (Fig. 1). The long-term trend of ocean heat is a major concern both in the scientific community and for the public at large. The increased temperatures cause a thermal expansion of water and a rise in sea level (Nerem et al., 2018; WCRP Global Sea Level Budget Group, 2018). The increase in ocean heat of  $(19.67 \pm 0.83) \times 10^{22}$  J in 2018 resulted in a 29.5 mm global mean sea level rise above the 1981–2010 average, and 1.4 mm above 2017. The resulting sea level rise exposes coastal freshwater supplies to saltwater intrusion, makes communities

\* Corresponding author: Lijing CHENG

Email: chenglij@mail.iap.ac.cn



**Fig. 1.** Upper panel: Change in 0–2000 m OHC from 1958 to 2018. Each bar shows the annual mean relative to a 1981–2010 baseline (positive in red and negative in blue). The green error bar indicates the 95% confidence interval, and the black line is the monthly time series. Lower panel: Annual mean OHC anomaly for the upper 2000 m in 2018 relative to a 1981–2010 baseline. Units:  $10^9$  J m $^{-2}$ . Source: IAP ocean analysis.

**Table 1.** Top five warmest years in the ocean since 1958. The OHC values in the upper 2000 m are the anomalies (units: J) relative to the 1981–2010 average.

Rank	Year	OHC (J)
1	2018	$19.67 \pm 0.83 \times 10^{22}$
2	2017	$18.76 \pm 0.80 \times 10^{22}$
3	2015	$17.99 \pm 0.70 \times 10^{22}$
4	2016	$17.47 \pm 0.76 \times 10^{22}$
5	2014	$16.22 \pm 0.70 \times 10^{22}$

more susceptible to storm surges, and threatens coastal infrastructure. The current sea level and its associated pattern is expected to continue and accelerate in the near future (Fasullo et al., 2018).

The increase in ocean heat (Fig. 1) also affects the planet’s weather systems, by raising air temperatures and supplying more moisture; warmer air can hold more moisture at a rate of about  $7\% \text{ }^\circ\text{C}^{-1}$  (Trenberth, 2011). In turn, this leads to increases in the intensity of storms and heavy rains (Patricola and Wehner, 2018; Trenberth et al., 2018). In 2018, the world experienced a number of major tropical storms, some of which developed very rapidly and many caused death and destruction. These included hurricanes Florence and Michael in the Atlantic, and major typhoons Jebi, Maria, Manghut and Trami in the

Pacific. Typhoon Mangkhut was the planet's most intense storm of 2018, with winds of  $287 \text{ km h}^{-1}$ ; it caused huge damage in Hong Kong. Several of these storms caused major flooding, especially Florence in the Carolinas. Other storms also led to major flooding events in Japan in July, and Kerala in India this past summer. Several hurricanes passed or hit Hawaii (e.g., Olivia, Lane), with Lane dropping more than 50 inches (1270 mm) of rain on Hawaii. It was the second-largest rainfall total from a tropical cyclone in the United States since 1950 (after Harvey in 2017). Hurricanes and other storms are natural phenomena and they are affected by many other factors besides ocean changes, but conditions allowing for the formation of severe hurricanes are occurring more often because of the record high OHC, with increases in intensity, lifetime, size, and especially increases in heavy rainfall (Patricola and Wehner, 2018; Trenberth et al., 2018; Wu et al., 2018).

Other consequences of ocean warming include declining ocean oxygen (Schmidtko et al., 2017), bleaching and death of corals (Hughes et al., 2018), melting sea ice and ice shelves directly through bottom heating (WCRP Global Sea Level Budget Group, 2018), increasing marine heat waves (MHWs) (Oliver et al., 2018), and altered impacts of natural variability such as ENSO (Fasullo et al., 2018). These consequences are not fully but at least partly attributable to ocean warming (IPCC, 2018). For example, 99% of the coral reefs will bleach if the surface warming reaches  $2^\circ\text{C}$  in 21st century because of ocean warming and acidification (IPCC, 2018). In 2018, MHWs were recorded in the Northwest and Southwest Pacific and the Northwest and Southwest Atlantic Ocean from sea surface temperature records. MHWs are defined as periods of extreme warm sea surface temperature that persist for days to months (Frölicher et al., 2018), supported by higher than normal OHC. MHWs in Northeast China during July–August 2018 took a heavy toll on sea cucumber farms. In addition to ocean-related effects, there are other indirect effects of ocean heating, such as increased drought intensity, heatwaves, and risk of wildfire.

Ocean warming clearly indicates a global warming since the 1950s, and the consequences of warming oceans are also clear. This global warming is a consequence of greenhouse gas (GHG) trapping of heat radiation within the Earth's system. Owing to the longevity of carbon dioxide and other GHGs, plus the thermal inertia of the climate system (Le Quéré, 2018), mitigating changes and the risk of major socioeconomic consequences due to ocean and global warming hinges on taking steps to immediately reduce GHGs emissions (Gattuso et al., 2018; IPCC, 2018).

## Data and methods

The method used here, which was developed at the IAP, involves calculating ocean temperatures down to 2000 m using observations from various measurement devices (Abraham et al., 2013; Cheng et al., 2017). The data are available at <http://159.226.119.60/cheng/>. The in-situ temperature measurements that were input into the IAP calculation are available from the National Oceanic and Atmospheric Administration/National Center for Environmental Information (NOAA/NCEI) (Boyer et al., 2013). The Argo observing network was implemented in around 2005, which has significantly improved the ocean-measurement capability (Argo, 2000). The Argo data are collected and made freely available by the International Argo Program and the contributing national programs (<http://www.argo.ucsd.edu>; <http://argo.jcommops.org>). The Argo Program is part of the Global Ocean Observing System. Note that relatively little is known about the deep ocean below 2000 m because observations at these depths are sparse (Argo data are mainly available for the upper 2000 m). The information on Hurricane Mangkhut came from <https://www.aljazeera.com/news/2018/09/china-hongkong-brace-typhoon-mangkhut-toll-rises-28-180916054503320.html>. ENSO information can be found at [https://origin.cpc.ncep.noaa.gov/products/analysis\\_monitoring/ensostuff/ONI\\_v5.php](https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php).

**Acknowledgements.** The IAP analysis is supported by the National Key Research and Development Program of China (Grant Nos. 2017YFA0603202 and 2016YFC1401705). We appreciate Tim BOYER from NOAA/NCEI to provide the in-situ ocean observations from NOAA/NCEI that were used in this analysis. Author contributions: L. C. and J. Z. are responsible for the IAP analysis and drafted the initial manuscript, with subsequent help from J. A., K. E. T. Authors J.A., K. E. T., J. F., Z. B., F. Y., L. W., X. C. and X. S. provide continual and vital support to the IAP analysis and activities, and contributed to refining this manuscript.

## REFERENCES

- Abraham, J. P., and Coauthors, 2013: A review of global ocean temperature observations: Implications for ocean heat content estimates and climate change. *Rev. Geophys.*, **51**, 450–483, <https://doi.org/10.1002/rog.20022>.
- Argo, 2000: Argo float data and metadata from global data assembly Centre (Argo GDAC). SEANOE, <https://doi.org/10.17882/42182>.
- Boyer, T. P., and Coauthors, 2013: World ocean database 2013. NOAA Atlas NESDIS 72, 209 pp.
- Cheng, L. J., and J. Zhu, 2018: 2017 was the warmest year on record for the global ocean. *Adv. Atmos. Sci.*, **35**(3), 261–263, <https://doi.org/10.1007/s00376-018-8011-z>.
- Cheng, L. J., K. E. Trenberth, J. Fasullo, T. Boyer, J. Abraham, and J. Zhu, 2017: Improved estimates of ocean heat content from 1960 to 2015. *Science Advances*, **3**, e1601545, <https://doi.org/10.1126/sciadv.1601545>.
- Cheng, L., J. Abraham, Z. Hausfather, and K. E. Trenberth, 2019: How fast are the oceans warming? *Science*, **363**, 128–129, <https://doi.org/10.1126/science.aav7619>.
- Cheng, L. J., G. J. Wang, J. P. Abraham, and G. Huang, 2018: Decadal ocean heat redistribution since the late 1990s and its association

- with key climate modes. *Climate*, **6**, 91, <https://doi.org/10.3390/cli6040091>.
- Fasullo, J. T., and R. S. Nerem, 2018: Altimeter-era emergence of the patterns of forced sea-level rise in climate models and implications for the future. *Proceedings of the National Academy of Sciences of the United States of America*, **115**, 12 944–12 949, <https://doi.org/10.1073/pnas.1813233115>.
- Fasullo, J. T., B. L. Otto-Bliesner, and S. Stevenson, 2018: ENSO's changing influence on temperature, precipitation, and wildfire in a warming climate. *Geophys. Res. Lett.*, **45**, 9216–9225, <https://doi.org/10.1029/2018GL079022>.
- Frölicher, T. L., E. M. Fischer, and N. Gruber, 2018: Marine heatwaves under global warming. *Nature*, **560**, 360–364, <https://doi.org/10.1038/s41586-018-0383-9>.
- Gattuso, J. P., and Coauthors, 2018: Ocean solutions to address climate change and its effects on marine ecosystems. *Frontiers in Marine Science*, **5**, 337, <https://doi.org/10.3389/fmars.2018.00337>.
- Hughes, T. P., and Coauthors, 2018: Global warming transforms coral reef assemblages. *Nature*, **556**, 492–496, <https://doi.org/10.1038/s41586-018-0041-2>.
- IPCC, 2018: Summary for Policymakers, Masson-Delmotte et al., Eds., *Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*, World Meteorological Organization, Geneva, Switzerland, 32 pp.
- Le Quéré, C., and Coauthors, 2018: Global carbon budget 2018. *Earth System Science Data*, **10**, 2141–2194, <https://doi.org/10.5194/essd-10-2141-2018>.
- Li, Y. L., W. Q. Han, A. X. Hu, G. A. Meehl, and F. Wang, 2018: Multidecadal changes of the upper Indian ocean heat content during 1965–2016. *J. Climate*, **31**, 7863–7884, <https://doi.org/10.1175/JCLI-D-18-0116.1>.
- Nerem, R. S., B. D. Beckley, J. T. Fasullo, B. D. Hamlington, D. Masters, and G. T. Mitchum, 2018: Climate -change-driven accelerated sea-level rise detected in the altimeter era. *Proceedings of the National Academy of Sciences of the United States of America*, **115**(9), 2022–2025, <https://doi.org/10.1073/pnas.1717312115>.
- Oliver, E. C. J., and Coauthors, 2018: Longer and more frequent marine heatwaves over the past century. *Nature Communications*, **9**, 1324, <https://doi.org/10.1038/s41467-018-03732-9>.
- Patricola, C. M., and M. F. Wehner, 2018: Anthropogenic influences on major tropical cyclone events. *Nature*, **563**, 339–346, <https://doi.org/10.1038/s41586-018-0673-2>.
- Rhein, M., and Coauthors, 2013: Observations: Ocean. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, T. F. Stocker et al., Eds., Cambridge University Press, Cambridge, UK, New York, NY, USA.
- Schmidtko, S., L. Stramma, and M. Visbeck, 2017: Decline in global oceanic oxygen content during the past five decades. *Nature*, **542**, 335–339, <https://doi.org/10.1038/nature21399>.
- Shi, J. R., S. P. Xie, and L. D. Talley, 2018: Evolving relative importance of the Southern Ocean and north Atlantic in anthropogenic ocean heat uptake. *J. Climate*, **31**, 7459–7479, <https://doi.org/10.1175/JCLI-D-18-0170.1>.
- Swart, N. C., S. T. Gille, J. C. Fyfe, and N. P. Gillett, 2018: Recent Southern Ocean warming and freshening driven by greenhouse gas emissions and ozone depletion. *Nature Geoscience*, **11**, 836–841, <https://doi.org/10.1038/s41561-018-0226-1>.
- Trenberth, K. E., 2011: Changes in precipitation with climate change. *Climate Research*, **47**, 123–138, <https://doi.org/10.3354/cr00953>.
- Trenberth, K. E., J. T. Fasullo, and M. A. Balmaseda, 2014: Earth's energy imbalance. *J. Climate*, **27**, 3129–3144, <https://doi.org/10.1175/JCLI-D-13-00294.1>.
- Trenberth, K. E., L. J. Cheng, P. Jacobs, Y. X. Zhang, and J. Fasullo, 2018: Hurricane Harvey links to ocean heat content and climate change adaptation. *Earth's Future*, **6**, 730–744, <https://doi.org/10.1029/2018EF000825>.
- von Schuckmann, K., and Coauthors, 2016: An imperative to monitor Earth's energy imbalance. *Nat. Clim. Change*, **6**, 138–144, <https://doi.org/10.1038/nclimate2876>.
- WCRP Global Sea Level Budget Group, 2018: Global sea-level budget 1993-present. *Earth System Science Data*, **10**(3), 1551–1590, <https://doi.org/10.5194/essd-10-1551-2018>.
- Wu, L. G., R. F. Wang, and X. F. Feng, 2018: Dominant role of the ocean mixed layer depth in the increased proportion of intense typhoons during 1980–2015. *Earth's Future*, **6**, 1518–1527, <https://doi.org/10.1029/2018EF000973>.